

Nonfactorizable Effects in Exclusive Charmless B Decays

Hai-Yang Cheng

Institute of Physics, Academia Sinica
Taipei, Taiwan 115, Republic of China

B. Tseng

Department of Physics, National Taiwan University
Taipei, Taiwan 106, Republic of China

Abstract

Nonfactorizable effects in charmless $B \rightarrow PP$, VP decays can be lumped into the effective parameters a_i that are linear combinations of Wilson coefficients, or equivalently absorbed into the effective number of colors N_c^{eff} . Naive factorization with $N_c^{\text{eff}} = 3$ fails to explain the CLEO data of $B^\pm \rightarrow \omega K^\pm$, indicating the first evidence for the importance of nonfactorizable contributions to the penguin amplitude. The decays $B^\pm \rightarrow \omega \pi^\pm$ dominated by tree amplitudes are sensitive to the interference between external and internal W -emission diagrams. Destructive interference implied by $N_c^{\text{eff}} = \infty$ leads to a prediction of $\mathcal{B}(B^\pm \rightarrow \omega \pi^\pm)$ which is about 2σ too small compared to experiment. Therefore, the CLEO data of $B^\pm \rightarrow \omega K^\pm$, $\omega \pi^\pm$ suffice to rule out $N_c^{\text{eff}} = 3$ and strongly disfavor $N_c^{\text{eff}} = \infty$ for rare charmless B decays. Factorization based on $N_c^{\text{eff}} \approx 2$ can accommodate the data of $B^\pm \rightarrow \omega K^\pm$, but it predicts a slightly smaller branching ratio of $B^\pm \rightarrow \omega \pi^\pm$ with $\mathcal{B}(B^\pm \rightarrow \omega \pi^\pm)/\mathcal{B}(B^\pm \rightarrow \omega K^\pm) = 0.6$. We briefly explain why the $1/N_c$ expansion is still applicable to the B meson decay once the correct large- N_c counting rule for the Wilson coefficient $c_2(m_b)$ is applied.

1. To describe the hadronic weak decays of mesons, the mesonic matrix elements are customarily evaluated under the factorization hypothesis so that they are factorized into the product of two matrix elements of single currents, governed by decay constants and form factors. In the naive factorization approach, the relevant Wilson coefficient functions for color-allowed external W -emission (or so-called “class-I”) and color-suppressed (class-II) internal W -emission amplitudes are given by $a_1 = c_1 + c_2/N_c$, $a_2 = c_2 + c_1/N_c$, respectively, with N_c the number of colors. In spite of its tremendous simplicity, naive factorization encounters two major difficulties. First, it never works for the decay rate of class-II decay modes, though it usually operates for class-I transition. For example, the predicted decay rate of the color-suppressed decay $D^0 \rightarrow \bar{K}^0 \pi^0$ in the naive approach is too small when compared with experiment (for a review, see [1]). Second, the hadronic matrix element under factorization is renormalization scale μ independent as the vector or axial-vector current is partially conserved. Consequently, the amplitude $c_i(\mu)\langle O \rangle_{\text{fact}}$ is not truly physical as the scale dependence of Wilson coefficients does not get compensation from the matrix elements. The first difficulty indicates that it is inevitable and mandatory to take into account nonfactorizable contributions, especially for class-II decays, to render the color suppression of internal W emission ineffective. In principle, the second difficulty also should not occur since the matrix elements of four-quark operators ought to be evaluated in the same renormalization scheme as that for Wilson coefficients and renormalized at the same scale μ .

Because there is only one single form factor (or Lorentz scalar) involved in the class-I or class II decay amplitude of $B(D) \rightarrow PP$, PV decays (P : pseudoscalar meson, V : vector meson), the effects of nonfactorization can be lumped into the effective parameters a_1 and a_2 [2]:¹

$$a_1^{\text{eff}} = c_1(\mu) + c_2(\mu) \left(\frac{1}{N_c} + \chi_1(\mu) \right), \quad a_2^{\text{eff}} = c_2(\mu) + c_1(\mu) \left(\frac{1}{N_c} + \chi_2(\mu) \right), \quad (1)$$

where nonfactorizable contributions are characterized by the parameters χ_1 and χ_2 . Taking the decay $B^- \rightarrow D^0 \pi^-$ as an example, we have [4, 5, 6]

$$\chi_1(\mu) = \varepsilon_8^{(BD,\pi)}(\mu) + \frac{a_1}{c_2} \varepsilon_1^{(BD,\pi)}(\mu), \quad \chi_2(\mu) = \varepsilon_8^{(B\pi,D)}(\mu) + \frac{a_2}{c_1} \varepsilon_1^{(B\pi,D)}(\mu), \quad (2)$$

where

$$\begin{aligned} \varepsilon_1^{(BD,\pi)}(\mu) &= \frac{\langle D^0 \pi^- | (\bar{d}u)_{V-A} (\bar{c}b)_{V-A} | B^- \rangle_{nf}}{\langle D^0 \pi^- | (\bar{d}u)_{V-A} (\bar{c}b)_{V-A} | B^- \rangle_f} = \frac{\langle D^0 \pi^- | (\bar{d}u)_{V-A} (\bar{c}b)_{V-A} | B^- \rangle}{\langle \pi^- | (\bar{d}u)_{V-A} | 0 \rangle \langle D^0 | (\bar{c}b)_{V-A} | B^- \rangle} - 1, \\ \varepsilon_8^{(BD,\pi)}(\mu) &= \frac{1}{2} \frac{\langle D^0 \pi^- | (\bar{d}\lambda^a u)_{V-A} (\bar{c}\lambda^a b)_{V-A} | B^- \rangle}{\langle \pi^- | (\bar{d}u)_{V-A} | 0 \rangle \langle D^0 | (\bar{c}b)_{V-A} | B^- \rangle}, \end{aligned} \quad (3)$$

are nonfactorizable terms originated from color-singlet and color-octet currents, respectively, $(\bar{q}_1 q_2)_{V-A} \equiv \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2$, and $(\bar{q}_1 \lambda^a q_2)_{V-A} \equiv \bar{q}_1 \lambda^a \gamma_\mu (1 - \gamma_5) q_2$. The subscript ‘f’ and

¹As pointed out in [3], the general amplitude of $B(D) \rightarrow VV$ decay consists of three independent Lorentz scalars, corresponding to S -, P - and D -wave amplitudes. Consequently, it is in general not possible to define an effective a_1 or a_2 unless nonfactorizable terms contribute in equal weight to all partial wave amplitudes.

‘nf’ in Eq. (3) stand for factorizable and nonfactorizable contributions, respectively, and the superscript (BD, π) in Eq. (2) means that the pion is factored out in the factorizable amplitude of $B \rightarrow D\pi$ and likewise for the superscript $(B\pi, D)$. In the large- N_c limit, $\varepsilon_1 = \mathcal{O}(1/N_c^2)$ and $\varepsilon_8 = \mathcal{O}(1/N_c)$ [6]. Therefore, the nonfactorizable term χ in the $N_c \rightarrow \infty$ limit is dominated by color-octet current operators. Since $|c_1/c_2| \gg 1$, it is evident from Eq. (1) that even a small amount of nonfactorizable contributions will have a significant effect on the color-suppressed class-II amplitude. Note that the effective parameters a_i^{eff} include all the contributions to the matrix elements and hence are μ independent [6]. If $\chi_{1,2}$ are universal (i.e. channel independent) in charm or bottom decays, then we still have a new factorization scheme in which the decay amplitude is expressed in terms of factorizable contributions multiplied by the universal effective parameters $a_{1,2}^{\text{eff}}$. The first systematical study of nonleptonic weak decays of heavy mesons within the framework of the improved factorization was carried out by Bauer, Stech, and Wirbel [7]. Phenomenological analyses of two-body decay data of D and B mesons indicate that while the generalized factorization hypothesis in general works reasonably well, the effective parameters $a_{1,2}^{\text{eff}}$ do show some variation from channel to channel, especially for the weak decays of charmed mesons [2, 5, 8]. An eminent feature emerged from the data analysis is that a_2^{eff} is negative in charm decay, whereas it becomes positive in bottom decay [2, 9, 6]:

$$a_2^{\text{eff}}(D \rightarrow \bar{K}\pi) \sim -0.50, \quad a_2^{\text{eff}}(B \rightarrow D\pi) \sim 0.26. \quad (4)$$

It should be stressed that since the magnitude of $a_{1,2}$ depends on the model results for form factors, the above values of a_2 should be considered as representative ones. The sign of a_2^{eff} is fixed by the observed destructive interference in $D^+ \rightarrow \bar{K}^0\pi^+$ and constructive interference in $B^- \rightarrow D^0\pi^-$. Eq. (4) then leads to

$$\chi_2(\mu \sim m_c; D \rightarrow \bar{K}\pi) \sim -0.36, \quad \chi_2(\mu \sim m_b; B \rightarrow D\pi) \sim 0.11. \quad (5)$$

In general the determination of χ_2 is easier and more reliable than χ_1 . The observation $|\chi_2(B)| \ll |\chi_2(D)|$ is consistent with the intuitive picture that soft gluon effects become stronger when the final-state particles move slower, allowing more time for significant final-state interactions after hadronization [2].

Phenomenologically, it is often to treat the number of colors N_c as a free parameter and fit it to the data. Theoretically, this amounts to defining an effective number of colors by

$$1/N_c^{\text{eff}} \equiv (1/N_c) + \chi. \quad (6)$$

It is clear from Eq. (5) that

$$N_c^{\text{eff}}(D \rightarrow \bar{K}\pi) \gg 3, \quad N_c^{\text{eff}}(B \rightarrow D\pi) \sim 2. \quad (7)$$

Consequently, the empirical rule of discarding subleading $1/N_c$ terms formulated in the large- N_c approach [10] is justified for exclusive charm decay; the dynamical origin of the $1/N_c$ expansion comes from the fact that the Fierz $1/N_c$ terms are largely compensated by

nonfactorizable effects in charm decay. Since the large- N_c approach implies $a_2^{\text{eff}} \sim c_2$ and since a_2^{eff} is observed to be positive in $B^- \rightarrow D^{(*)}\pi^-(\rho^-)$ decays, one may wonder why is the $1/N_c$ expansion no longer applicable to the B meson? Contrary to the common belief, a careful study shows this is not the case. As pointed out in [6], the large- N_c color counting rule for the Wilson coefficient $c_2(\mu)$ is different at $\mu \sim m_b$ and $\mu \sim m_c$ due to the presence of the large logarithm at $\mu \sim m_c$. More specifically, $c_2(m_b) = \mathcal{O}(1/N_c)$ and $c_2(m_c) = \mathcal{O}(1)$. Recalling that $c_1 = \mathcal{O}(1)$, it follows that in the large- N_c limit [6]:

$$a_2^{\text{eff}} = \begin{cases} c_2(m_c) + \mathcal{O}(1/N_c) & \text{for the } D \text{ meson,} \\ c_2(m_b) + c_1(m_b) \left(\frac{1}{N_c} + \varepsilon_8(m_b) \right) + \mathcal{O}(1/N_c^3) & \text{for the } B \text{ meson.} \end{cases} \quad (8)$$

Therefore, *a priori* the $1/N_c$ expansion does not demand a negative a_2^{eff} for bottom decay! and $N_c^{\text{eff}}(B \rightarrow D\pi) \sim 2$ is not in conflict with the large- N_c approach! It should be remarked that although χ_2 is positive in two-body decays of the B meson, some theoretical argument suggests that it may become negative for high multiplicity decay modes [6].

Thus far the nonfactorization effect is discussed at the purely phenomenological level. It is thus important to have a theoretical estimate of χ_i even approximately. Unfortunately, all existing theoretical calculations based on the QCD sum rule [11], though confirm the cancellation between the $1/N_c$ Fierz terms and nonfactorizable soft gluon effects [12], tend to predict a negative χ in $\bar{B}^0 \rightarrow D^+\pi^-$, $D^0\pi^0$ and $B \rightarrow J/\psi K(K^*)$ decays. This tantalizing issue should be clarified and resolved in the near future. It is interesting to remark that, relying on a different approach, namely, the three-scale PQCD factorization theorem, to tackle the nonfactorization effect, one of us and Li [13] are able to explain the sign change of χ_2 from bottom to charm decays.

For B meson decay, the effective parameters $a_{1,2}^{\text{eff}}$ have been determined so far only for $B \rightarrow D^{(*)}\pi(\rho)$ and $B \rightarrow J/\psi K^{(*)}$ where nonfactorizable effects amount to having $N_c^{\text{eff}} \sim 2$. Recently, several exclusive charmless rare B decay modes have been reported for the first time by CLEO [14] and many of them are dominated by the penguin mechanism. It is thus important to know (i) does the constructive interference of tree amplitudes persist in class-III charmless B decay? (class-III transitions receive contributions from both external and internal W emissions), and (ii) is $N_c^{\text{eff}} \sim 2$ still applicable to the penguin amplitude? Whether $N_c^{\text{eff}} \sim 2$ or $N_c^{\text{eff}} \sim \infty$ for B decay to two light mesons is still under debate. For example, predictions for exclusive charmless B decay are presented in [15] for $N_c^{\text{eff}} = \infty$. Recently, it was argued in [16] that the CLEO data of two-body charmless B decays can be accommodated by $0 \leq 1/N_c^{\text{eff}} \leq 0.5$ with $N_c^{\text{eff}} = \infty$ being more preferred. In this Letter we shall demonstrate that naive factorization (i.e. $N_c^{\text{eff}} = 3$) is ruled out by the CLEO data of $B^\pm \rightarrow \omega K^\pm$ and $N_c^{\text{eff}} = \infty$ is strongly disfavored by the data of $B^\pm \rightarrow \omega \pi^\pm$. This implies the applicability of $N_c^{\text{eff}} \sim 2$ to the rare charmless B decays.

2. In this section we will consider the decay modes dominated by penguin diagrams in order to study their N_c^{eff} dependence. It was pointed out in [17] that the parameters a_2 , a_3 and a_5 are strongly dependent on N_c^{eff} and the rates dominated by these coefficients can have large variation. For example, the decay widths of $B^- \rightarrow \omega K^{(*)-}$, $B^0 \rightarrow \omega K^0$, ρK^{*0} ,

$B_s \rightarrow \eta\omega, \eta\phi, \omega\phi, \dots$, etc. have strong N_c dependence [17]. We shall see that the branching ratio of $B^- \rightarrow \omega K^-$ has its lowest value near $N_c^{\text{eff}} \sim 3 - 4$ and hence the naive factorization with $N_c^{\text{eff}} = 3$ is ruled out by experiment.

Before proceeding we briefly sketch the calculational framework. The relevant effective $\Delta B = 1$ weak Hamiltonian is

$$\mathcal{H}_{\text{eff}}(\Delta B = 1) = \frac{G_F}{\sqrt{2}} [V_{ub}V_{uq}^*(c_1 O_1^u + c_2 O_2^u) + V_{cb}V_{cq}^*(c_1 O_1^c + c_2 O_2^c) - V_{tb}V_{tq}^* \sum_{i=3}^{10} c_i O_i] + \text{h.c.}, \quad (9)$$

where $q = u, d, s$, and

$$\begin{aligned} O_1^u &= (\bar{u}b)_{V-A}(\bar{q}u)_{V-A}, & O_2^u &= (\bar{q}b)_{V-A}(\bar{u}u)_{V-A}, \\ O_{3(5)} &= (\bar{q}b)_{V-A} \sum_{q'} (\bar{q}'q')_{V-A(V+A)}, & O_{4(6)} &= (\bar{q}_\alpha b_\beta)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\alpha)_{V-A(V+A)}, \\ O_{7(9)} &= \frac{3}{2}(\bar{q}b)_{V-A} \sum_{q'} e_{q'}(\bar{q}'q')_{V+A(V-A)}, & O_{8(10)} &= \frac{3}{2}(\bar{q}_\alpha b_\beta)_{V-A} \sum_{q'} e_{q'}(\bar{q}'_\beta q'_\alpha)_{V+A(V-A)}, \end{aligned} \quad (10)$$

where O_3 - O_6 are QCD penguin operators and O_7 - O_{10} originate from electroweak penguin diagrams. As noted in passing, in order to ensure the renormalization-scale and -scheme independence for the physical amplitude, the matrix elements of 4-quark operators have to be evaluated in the same renormalization scheme as that for Wilson coefficients and renormalized at the same scale μ . Before utilizing factorization, it is necessary to take into account QCD and electroweak corrections to matrix elements:

$$\langle O_i(\mu) \rangle = \left[\mathbf{I} + \frac{\alpha_s(\mu)}{4\pi} \hat{m}_s(\mu) + \frac{\alpha}{4\pi} \hat{m}_e(\mu) \right]_{ij} \langle O_j^{\text{tree}} \rangle, \quad (11)$$

so that $c_i(\mu) \langle O_i(\mu) \rangle = \tilde{c}_i \langle O_i^{\text{tree}} \rangle$, where

$$\tilde{c}_i = \left[\mathbf{I} + \frac{\alpha_s(\mu)}{4\pi} \hat{m}_s(\mu) + \frac{\alpha}{4\pi} \hat{m}_e(\mu) \right]_{ji} c_j(\mu). \quad (12)$$

Then the factorization approximation is applied to the hadronic matrix elements of the tree operator O^{tree} . Perturbative QCD and electroweak corrections to the matrices $\hat{m}_s$ and $\hat{m}_e$ have been calculated in [18, 19, 15, 16]. Using the next-to-leading order $\Delta B = 1$ Wilson coefficients obtained in the 't Hooft-Veltman scheme and the naive dimension regularization scheme at $\mu = 4.4$ GeV, $\Lambda_{\overline{\text{MS}}}^{(5)} = 225$ MeV and $m_t = 170$ GeV in Table 22 of [20], we obtain the renormalization-scheme and -scale independent Wilson coefficients $\tilde{c}_i$ at $k^2 = m_b^2/2$:²

$$\begin{aligned} \tilde{c}_1 &= 1.187, & \tilde{c}_2 &= -0.312, \\ \tilde{c}_3 &= 0.0236 + i0.0048, & \tilde{c}_4 &= -0.0547 - i0.0143, \\ \tilde{c}_5 &= 0.0164 + i0.0048, & \tilde{c}_6 &= -0.0640 - i0.0143, \\ \tilde{c}_7 &= -(0.0757 + i0.0558)\alpha, & \tilde{c}_8 &= 0.057\alpha, \\ \tilde{c}_9 &= -(1.3648 + i0.0558)\alpha, & \tilde{c}_{10} &= 0.264\alpha. \end{aligned} \quad (13)$$

²The values of $\tilde{c}_i$ given in (13) are slightly different from that shown in [21]. Here we have included QCD vertex corrections to $\tilde{c}_1, \dots, \tilde{c}_6$ (see [16] for the expressions of vertex corrections). It should be stressed that long-distance nonfactorizable effects have not been included so far.

We now apply the Hamiltonian (9) and factorization to the decay $B^- \rightarrow \omega K^-$ and obtain

$$\begin{aligned}
A(B^- \rightarrow \omega K^-) &= \frac{G_F}{\sqrt{2}} \left\{ V_{ub}V_{us}^* (a_1 X_1 + a_2 X_{2u} + a_1 X_3) \right. \\
&\quad - V_{tb}V_{ts}^* \left[\left(a_4 + a_{10} - 2(a_6 + a_8) \frac{m_K^2}{(m_s + m_u)(m_b + m_u)} \right) X_1 \right. \\
&\quad + \frac{1}{2} (4a_3 + 4a_5 + a_7 + a_9) X_{2u} \\
&\quad \left. \left. + \left(a_4 + a_{10} - 2(a_6 + a_8) \frac{m_B^2}{(m_s + m_u)(m_b + m_u)} \right) X_3 \right] \right\}, \quad (14)
\end{aligned}$$

where $a_{2i} \equiv \tilde{c}_{2i} + \frac{1}{N_c} \tilde{c}_{2i-1}$, $a_{2i-1} \equiv \tilde{c}_{2i-1} + \frac{1}{N_c} \tilde{c}_{2i}$, and X_i are factorizable terms:

$$\begin{aligned}
X_1 &\equiv \langle K^- | (\bar{s}u)_{V-A} | 0 \rangle \langle \omega | (\bar{u}b)_{V-A} | B^- \rangle = -i\sqrt{2} f_K m_\omega A_0^{B\omega}(m_K^2)(\varepsilon \cdot p_B), \\
X_{2q} &\equiv \langle \omega | (\bar{q}q)_{V-A} | 0 \rangle \langle K^- | (\bar{s}b)_{V-A} | B^- \rangle = -i\sqrt{2} f_\omega m_\omega F_1^{BK}(m_\omega^2)(\varepsilon \cdot p_B), \\
X_3 &\equiv \langle \omega K^- | (\bar{s}u)_{V-A} | 0 \rangle \langle 0 | (\bar{u}b)_{V-A} | B^- \rangle, \quad (15)
\end{aligned}$$

with ε the polarization vector of the ω meson, and A_1 , F_1 the form factors defined in [22]. Just as in the case of tree amplitudes, one can show that nonfactorizable effects in the penguin amplitudes of $B \rightarrow PP$, VP decays can be absorbed into the effective penguin coefficients. This amounts to replacing N_c in the penguin coefficients a_i ($i = 3, \dots, 10$) by $(N_c^{\text{eff}})_i$. (It must be emphasized that the factor of N_c appearing in any place other than a_i should *not* be replaced by N_c^{eff} .) For simplicity, we will assume $(N_c^{\text{eff}})_1 \approx (N_c^{\text{eff}})_2 \dots \approx (N_c^{\text{eff}})_{10}$ so that the subscript i can be dropped. For $N_c^{\text{eff}} = 3$, the QCD-penguin Wilson coefficients are numerically given by $\text{Re } a_3 = 0.0054$, $\text{Re } a_4 = -0.0468$, $\text{Re } a_5 = -0.0049$, and $\text{Re } a_6 = -0.0585$. From Eq. (14) we see that a large cancellation occurs in the QCD penguin amplitude due to the large compensation between a_3 and a_5 , a_4 and a_6 . Since $|V_{tb}V_{ts}^*| \gg |V_{ub}V_{us}^*|$, the decay rate of $B^\pm \rightarrow \omega K^\pm$ has its minimum around $N_c^{\text{eff}} \sim 3 - 4$ (see Fig. 1), as noticed in [17, 15] and analyzed in detail in [16].

Neglecting the W -annihilation contribution denoted by X_3 , and using $f_K = 160$ MeV, $f_\omega = 195$ MeV for decay constants, $A_0^{B\omega}(0) = 0.28$, $F_1^{BK}(0) = 0.34$ [22], and dipole q^2 dependence for form factors A_0 and F_1 [23], $m_u = 5$ MeV, $m_d = 10$ MeV, $m_s = 175$ MeV, $m_b = 5$ GeV for quark masses, $\tau(B^\pm) = (1.66 \pm 0.04)$ ps [24] for the charged B lifetime, and $A = 0.804$, $\lambda = 0.22$, $\eta = 0.30$, $\rho = \pm 0.30$ [21] for Wolfenstein parameters [25], we obtain the averaged branching ratios of $B^\pm \rightarrow \omega K^\pm$ defined by

$$\mathcal{B}(B^\pm \rightarrow \omega K^\pm) \equiv \frac{1}{2} \left[\mathcal{B}(B^+ \rightarrow \omega K^+) + \mathcal{B}(B^- \rightarrow \omega K^-) \right], \quad (16)$$

in Table I and Fig. 1. We see that the prediction $\mathcal{B}(B^\pm \rightarrow \omega K^\pm) = 1.44 \times 10^{-6}$ at $N_c^{\text{eff}} = 3$ and $\rho = -0.30$ is off by 2σ from the experimental result [14]

$$\mathcal{B}(B^\pm \rightarrow \omega K^\pm) = (1.2_{-0.5}^{+0.7} \pm 0.2) \times 10^{-5}. \quad (17)$$

This shows that naive factorization with $N_c^{\text{eff}} = 3$ (or $\chi = 0$) fails to explain the decay rate of $B^\pm \rightarrow \omega K^\pm$. It is clear from Table I or Fig. 1 that, for $\rho < 0$, $N_c^{\text{eff}} = 2$ is slightly better than

Table I. Averaged branching ratios for charmless B decays, where “Tree” refers to branching ratios from tree diagrams only, “Tree+QCD” from tree and QCD penguin diagrams, and “Tree+QCD+QED” from tree, QCD and electroweak (EW) penguin diagrams. Predictions are made for $k^2 = m_b^2/2$, $\eta = 0.30$, $\rho = 0.30$ (the first number in parentheses) and $\rho = -0.30$ (the second number in parentheses).

Decay	N_c^{eff}	Tree	Tree+QCD	Tree+QCD+EW	Exp. [14]
$B^\pm \rightarrow \omega K^\pm$	2	6.53×10^{-7}	$(2.43, 6.82) 10^{-6}$	$(3.28, 8.31) 10^{-6}$	$(1.2_{-0.5}^{+0.7} \pm 0.2) \times 10^{-5}$
	3	4.49×10^{-7}	$(2.58, 9.63) 10^{-7}$	$(0.27, 1.44) 10^{-6}$	
	∞	1.56×10^{-7}	$(9.49, 6.44) 10^{-6}$	$(8.46, 5.61) 10^{-6}$	
$\bar{B}^0 \rightarrow \omega \bar{K}^0$	2	5.25×10^{-8}	$(3.24, 4.46) 10^{-6}$	$(3.78, 5.09) 10^{-6}$	
	3	4.64×10^{-9}	$(1.37, 2.10) 10^{-7}$	$(2.89, 3.96) 10^{-7}$	
	∞	6.45×10^{-8}	$(6.28, 8.16) 10^{-6}$	$(5.18, 6.89) 10^{-6}$	
$B^\pm \rightarrow \omega \pi^\pm$	2	9.71×10^{-6}	$(1.15, 0.53) 10^{-5}$	$(1.16, 0.52) 10^{-5}$	$(1.2_{-0.5}^{+0.7} \pm 0.2) \times 10^{-5}$
	3	6.23×10^{-6}	$(7.07, 3.93) 10^{-6}$	$(7.16, 3.76) 10^{-6}$	
	∞	1.57×10^{-6}	$(1.49, 1.93) 10^{-6}$	$(1.52, 1.78) 10^{-6}$	

$N_c^{\text{eff}} = \infty$ and yields a branching ratio of order 8.3×10^{-6} , in agreement with experiment.³ Note that the electroweak penguin contribution is constructive at $N_c^{\text{eff}} = 2$ and destructive at $N_c^{\text{eff}} = \infty$. For $\rho > 0$, $N_c^{\text{eff}} = \infty$ works better than $N_c^{\text{eff}} = 2$ and can accommodate the data. However, we have shown in [21] that a positive ρ is quite disfavored by data as it predicts $\mathcal{B}(B^\pm \rightarrow \eta \pi^\pm) \sim 1 \times 10^{-5}$, which is marginally larger than the CLEO measured upper limit: 0.8×10^{-5} [14].

A similar strong N_c^{eff} dependence also can be observed in $B_d^0 \rightarrow \omega K^0$ decay.⁴ The expression of its factorizable amplitude is simpler than the charged B meson:

$$\begin{aligned}
A(B_d^0 \rightarrow \omega K^0) = & \frac{G_F}{\sqrt{2}} \left\{ V_{ub} V_{us}^* a_2 X_{2u} - V_{tb} V_{ts}^* \left[\frac{1}{2} (4a_3 + 4a_5 + a_7 + a_9) X_{2u} \right. \right. \\
& \left. \left. + \left(a_4 - \frac{1}{2} a_{10} - (2a_6 - a_8) \frac{m_K^2}{(m_s + m_d)(m_b + m_d)} \right) X_1 \right] \right\}. \quad (18)
\end{aligned}$$

The averaged decay rate $\Gamma(\bar{B}^0 \rightarrow \omega \bar{K}^0) \equiv \frac{1}{2} [\Gamma(B^0 \rightarrow \omega K^0) + \Gamma(\bar{B}^0 \rightarrow \omega \bar{K}^0)]$ is minimal near $N_c^{\text{eff}} \sim 4$ (see Fig. 2). The branching ratio predicted by $N_c^{\text{eff}} = \infty$ is slightly larger than that by $N_c^{\text{eff}} = 2$ (see also Table I).

3. In the last section we have demonstrated that $N_c^{\text{eff}} = 3$ is ruled out as it fails to describe the decays $B^\pm \rightarrow \omega K^\pm$. This means that it is mandatory to take into account the nonfactorizable effect in the charmless B decay amplitude. However, it is not easy to discern

³Our conclusions for $N_c^{\text{eff}} = 2$ and $N_c^{\text{eff}} = \infty$ are different from that in [16] which claimed that a value of $1/N_c^{\text{eff}}$ in the range $0.15 \leq 1/N_c^{\text{eff}} \leq 0.55$ is disfavored by the data of $B^\pm \rightarrow \omega K^\pm$ and that $0 \leq 1/N_c^{\text{eff}} \leq 0.15$ is preferred.

⁴As noted before, nonfactorizable effects in $B \rightarrow VV$ generally cannot be absorbed into the effective parameters a_i . Hence, we will not discuss $B \rightarrow \omega K^*$, ρK^* decays.

between $N_c^{\text{eff}} = \infty$ and $N_c^{\text{eff}} = 2$ from $B \rightarrow \omega K$ decays, though the latter can explain the data of $B^\pm \rightarrow \omega K^\pm$ and is more preferred. It is thus important to have a more decisive test on N_c^{eff} . For this purpose, we shall focus in this section the decay modes dominated by the tree diagrams and sensitive to the interference between external and internal W -emission amplitudes. The fact that $N_c^{\text{eff}} = 2$ ($N_c^{\text{eff}} = \infty$) implies constructive (destructive) interference will enable us to differentiate between them. Good examples are the class-III modes: $B^\pm \rightarrow \pi^0 \pi^\pm$, $\eta \pi^\pm$, $\pi^0 \rho^\pm$, $\omega \pi^\pm$, $\dots$. Since $B^- \rightarrow \omega \pi^-$ is the only channel in the list that has been measured by CLEO, we will first investigate this mode.

Under factorization, the decay amplitude of $B^- \rightarrow \omega \pi^-$ is given by

$$\begin{aligned}
A(B^- \rightarrow \omega \pi^-) = & \frac{G_F}{\sqrt{2}} \left\{ V_{ub} V_{ud}^* (a_1 X'_1 + a_2 X'_{2u} + a_1 X'_3) \right. \\
& - V_{tb} V_{td}^* \left[\left(a_4 + a_{10} - 2(a_6 + a_8) \frac{m_\pi^2}{(m_b + m_u)(m_u + m_d)} \right) X'_1 \right. \\
& + \frac{1}{2} (4a_3 + 2a_4 + 4a_5 + a_7 + a_9 - a_{10}) X'_{2u} \\
& \left. \left. + \left(a_4 + a_{10} - 2(a_6 + a_8) \frac{m_B^2}{(m_b + m_u)(m_u + m_d)} \right) X'_3 \right] \right\}, \quad (19)
\end{aligned}$$

with the expressions of X'_i similar to (15). Since

$$V_{ub} V_{ud}^* = A\lambda^3(\rho - i\eta), \quad V_{cb} V_{cd}^* = -A\lambda^3, \quad V_{tb} V_{td}^* = A\lambda^3(1 - \rho + i\eta), \quad (20)$$

in terms of the Wolfenstein parametrization [25], are of the same order of magnitude, it is clear that $B^- \rightarrow \omega \pi^-$ is dominated by external and internal W emissions and that penguin contributions are suppressed by the smallness of penguin coefficients. In the limit of $N_c^{\text{eff}} \rightarrow \infty$, we have $a_1 = c_1$ and $a_2 = c_2$, which in turn imply a destructive interference of tree amplitudes in $B^\pm \rightarrow \omega \pi^\pm$. It is easily seen from Eq. (13) that the interference becomes constructive when $N_c^{\text{eff}} < 3.8$. From Fig. 3 or Table I we see that the averaged branching ratio of $B^\pm \rightarrow \omega \pi^\pm$ has its lowest value of order 2×10^{-6} at $N_c^{\text{eff}} = \infty$ and then increases with $1/N_c^{\text{eff}}$. Since experimentally [14]

$$\mathcal{B}(B^\pm \rightarrow \omega \pi^\pm) = (1.2_{-0.5}^{+0.7} \pm 0.2) \times 10^{-5}, \quad (21)$$

it is evident that $N_c^{\text{eff}} = \infty$ is strongly disfavored by the data. Note that though the predicted branching ratio $\mathcal{B}(B^\pm \rightarrow \omega \pi^\pm) = 1.16 \times 10^{-5}$ for $N_c^{\text{eff}} = 2$ and $\rho = 0.30$ (see Table I) is in good agreement with experiment, we have discussed in passing that a positive ρ seems to be ruled out [21]. For $\rho < 0$, our prediction $\mathcal{B}(B^\pm \rightarrow \omega \pi^\pm) = 0.52 \times 10^{-5}$ is on the verge of the lower side of the CLEO data. Of course, the CLEO measurement can be more satisfactorily explained by having a much smaller N_c^{eff} , but this possibility is very unlikely as it implies a large nonfactorization effect in $B^\pm \rightarrow \omega \pi^\pm$. Recalling that the magnitude of nonfactorizable term is $\chi \sim 0.1$ in $B \rightarrow D\pi$ decay and that the energy release in the process $B \rightarrow \omega \pi$ is larger than that in $B \rightarrow D\pi$, it is thus expected physically that $\chi \lesssim 0.1$ for the former.

Using $N_c^{\text{eff}} = 2$, we find a slightly smaller branching ratio for $B^\pm \rightarrow \omega\pi^\pm$ with

$$\frac{\mathcal{B}(B^\pm \rightarrow \omega\pi^\pm)}{\mathcal{B}(B^\pm \rightarrow \omega K^\pm)} = 0.61. \quad (22)$$

Since theoretically it is difficult to see how the branching ratio of $B^\pm \rightarrow \omega\pi^\pm$ can be enhanced from 0.5×10^{-5} to 1.2×10^{-5} , it is thus important to have a refined and improved measurement of this decay mode.

In analogue to the decays $B \rightarrow D^{(*)}\pi(\rho)$, the interference effect of tree amplitudes in class-III charmless B decay can be tested by measuring the ratios:

$$R_1 \equiv 2 \frac{\mathcal{B}(B^- \rightarrow \pi^- \pi^0)}{\mathcal{B}(\bar{B}^0 \rightarrow \pi^- \pi^+)}, \quad R_2 \equiv 2 \frac{\mathcal{B}(B^- \rightarrow \rho^- \pi^0)}{\mathcal{B}(\bar{B}^0 \rightarrow \rho^- \pi^+)}, \quad R_3 \equiv 2 \frac{\mathcal{B}(B^- \rightarrow \pi^- \rho^0)}{\mathcal{B}(\bar{B}^0 \rightarrow \pi^- \rho^+)}. \quad (23)$$

Since penguin contributions are very small, to a good approximation we have

$$\begin{aligned} R_1 &= \frac{\tau(B^-)}{\tau(B_d^0)} \left(1 + \frac{a_2}{a_1}\right)^2, \\ R_2 &= \frac{\tau(B^-)}{\tau(B_d^0)} \left(1 + \frac{f_\pi}{f_\rho} \frac{A_0^{B\rho}(m_\pi^2)}{F_1^{B\pi}(m_\rho^2)} \frac{a_2}{a_1}\right)^2, \\ R_3 &= \frac{\tau(B^-)}{\tau(B_d^0)} \left(1 + \frac{f_\rho}{f_\pi} \frac{F_1^{B\pi}(m_\rho^2)}{A_0^{B\rho}(m_\pi^2)} \frac{a_2}{a_1}\right)^2. \end{aligned} \quad (24)$$

Evidently, the ratios R_i are greater (less) than unity when the interference is constructive (destructive). Numerically we find

$$R_1 = \begin{cases} 1.74, \\ 0.58, \end{cases} \quad R_2 = \begin{cases} 1.40, \\ 0.80, \end{cases} \quad R_3 = \begin{cases} 2.50 & \text{for } N_c^{\text{eff}} = 2, \\ 0.26 & \text{for } N_c^{\text{eff}} = \infty, \end{cases} \quad (25)$$

where use of $\tau(B_d^0) = (1.55 \pm 0.04)$ ps [24], $f_\rho = 216$ MeV, $A_0^{B\rho}(0) = 0.28$ [23] has been made. Hence, a measurement of R_i (in particular R_3), which has the advantage of being independent of the parameters ρ and η , will constitute a very useful test on the effective number of colors N_c^{eff} .

We would like to stress once again that the observation that $N_c^{\text{eff}} = \infty$ is very likely to be ruled out in charmless B decay does *not* imply the inapplicability of the large- N_c approach to the B meson case. As explained before, the correct large- N_c counting rule for the Wilson coefficient $c_2(m_b)$ is proportional to $1/N_c$. Consequently, a nontrivial a_2^{eff} , given by $c_2(m_b) + \frac{1}{N_c^{\text{eff}}} c_1(m_b)$, starts at the order of $1/N_c$ and hence N_c^{eff} cannot go to infinity.

4. To conclude, by absorbing the nonfactorizable effects into the effective number of colors N_c^{eff} , we have shown that $N_c^{\text{eff}} = 3$ is ruled out by the CLEO data of $B^\pm \rightarrow \omega K^\pm$, implying the inapplicability of naive factorization to charmless B decays, and that $N_c^{\text{eff}} = \infty$ is strongly disfavored by the experimental measurement of $B^\pm \rightarrow \omega\pi^\pm$, indicating a constructive interference in class-III charmless B decays.

Since the energy release in charmless two-body decays of the B meson is generally slightly larger than that in $B \rightarrow D^{(*)}\pi$, $D^{(*)}\rho$, it is natural to expect that N_c^{eff} for the B decay into two light mesons is close to $N_c^{\text{eff}}(B \rightarrow D\pi) \approx 2$. We have shown that $N_c^{\text{eff}} \approx 2$ can accommodate the data of $B^\pm \rightarrow \omega K^\pm$, but it predicts a slightly smaller branching ratio of $B^\pm \rightarrow \omega \pi^\pm$ with $\mathcal{B}(B^\pm \rightarrow \omega \pi^\pm)/\mathcal{B}(B^\pm \rightarrow \omega K^\pm) = 0.6$. Thus far, the discussion of nonfactorization in rare B decay is at the purely phenomenological level. It remains to be a challenge, especially in the framework of the QCD sum rule, to compute it theoretically.

ACKNOWLEDGMENT: This work was supported in part by the National Science Council of ROC under Contract Nos. NSC86-2112-M-001-020 and NSC86-2112-M-001-010-Y.

References

- [1] H.Y. Cheng, *Int. J. Mod. Phys. A***4**, 495 (1989).
- [2] H.Y. Cheng, *Phys. Lett. B***395**, 345 (1994); in *Particle Theory and Phenomenology*, XVII International Karimierz Meeting on Particle Physics, Iowa State University, May 1995, edited by K.E. Lassila *et al.* (World Scientific, Singapore, 1996), p.122 .
- [3] A.N. Kamal and A.B. Santra, Alberta Thy-31-94 (1994); *Z. Phys. C***72**, 91 (1996).
- [4] J.M. Soares, *Phys. Rev. D***51**, 3518 (1995);
- [5] A.N. Kamal, A.B. Santra, T. Uppal, and R.C. Verma, *Phys. Rev. D***53**, 2506 (1996).
- [6] M. Neubert and B. Stech, CERN-TH/97-99 [hep-ph/9705292], to appear in *Heavy Flavours*, edited by A.J. Buras and M. Lindner, 2nd ed. (World Scientific, Singapore).
- [7] M. Bauer, B. Stech, and M. Wirbel, *Z. Phys. C***34**, 103 (1987).
- [8] H.Y. Cheng, *Z. Phys. C***69**, 647 (1996).
- [9] H.Y. Cheng and B. Tseng, *Phys. Rev. D***51**, 6295 (1995).
- [10] A.J. Buras, J.-M. Gérard, and R. Rückl, *Nucl. Phys. B***268**, 16 (1986).
- [11] B. Blok and M. Shifman, *Nucl. Phys. B***389**, 534 (1993); A. Khodjamirian and R. Rückl, MPI-PhT/94-26 (1994); *Nucl. Phys. (Proc. Suppl.)* **39BC**, 396 (1995); I. Halperin, *Phys. Lett. B***349**, 548 (1995).
- [12] B. Blok and M. Shifman, *Sov. J. Nucl. Phys.* **45**, 35, 301, 522 (1987).
- [13] H.-n. Li and B. Tseng, NCKU-HEP-97-02 [hep-ph/9706441].

- [14] CLEO Collaboration, talks presented by B. Behrens and J. Alexander at *The Second International Conference on B Physics and CP Violation*, March 24-27, 1997, Honolulu, Hawaii; F. Würthwein, CALT-68-2121 [hep-ex/9706010]; P.A. Pomianowski, talk presented at *The Sixth Conference on the Intersections of Particle and Nuclear Physics*, Big Sky Montana, May 28, 1997.
- [15] G. Kramer, W.F. Palmer, and H. Simma, *Z. Phys.* **C66**, 429 (1995); *Nucl. Phys.* **B428**, 77 (1994).
- [16] A. Ali and C. Greub, DESY 97-126 [hep-ph/9707251].
- [17] A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, *Phys. Lett.* **B320**, 170 (1994).
- [18] A.J. Buras, M. Jamin, M.E. Lautenbacher, and P.H. Weisz, *Nucl. Phys.* **B370**, 69 (1992); A.J. Buras, M. Jamin, and M.E. Lautenbacher, *Nucl. Phys.* **B408**, 209 (1993).
- [19] R. Fleischer, *Z. Phys.* **C58**, 483 (1993).
- [20] G. Buchalla, A.J. Buras, and M.E. Lautenbacher, *Rev. Mod. Phys.* **68**, 1125 (1996).
- [21] H.Y. Cheng and B. Tseng, IP-ASTP-03-97 [hep-ph/9707316].
- [22] M. Wirbel, B. Stech, and M. Bauer, *Z. Phys.* **C29**, 637 (1985).
- [23] H.Y. Cheng, C.Y. Cheung, and C.W. Hwang, *Phys. Rev.* **D55**, 1559 (1997).
- [24] J.D. Richman, hep-ex/9701014.
- [25] L. Wolfenstein, *Phys. Rev. Lett.* **13**, 562 (1984).




Figure 1: The branching ratio of $B^\pm \rightarrow \omega K^\pm$ vs $1/N_c^{\text{eff}}$. The solid and dashed curves are for $\rho = -0.30$ and $\rho = 0.30$ respectively.




Figure 2: Same as Fig. 1 except for $B^{(\pm)0} \rightarrow \omega K^{(\pm)0}$.




Figure 3: Same as Fig. 1 except for $B^\pm \rightarrow \omega \pi^\pm$.